

Cutaneous Lymphoma Subsection of the SSMDT

Primary Cutaneous Lymphomas are a heterogeneous and rare group of disorders that may be treated by consultant dermatologists with skin-directed therapies but sometimes require specialist input from clinical oncologists and haematologists. Clinical-pathological correlation is often crucial for diagnosis in this group.

The cutaneous lymphoma sub-MDT is part of the wider Skin Cancer SSMDT based in Leicester. However, due to the rarity of this group of cutaneous malignancies, and significant differences in treatment modalities in comparison to the majority of skin cancers, these patients are discussed separately with a different group of specialists giving advice to the MDT.

To refer a patient to the cutaneous lymphoma sub-MDT, please see below:

Who?	All new diagnoses of primary cutaneous lymphoma (see Table 1 below for categories) Patients refractory to skin-directed therapy Patients where TSEBT or ECP is being considered
How?	Email CancerCentreSkin@UHL-tr.nhs.uk and C.C. co-ordinator Bansal.Deol@UHL-tr.nhs.uk
What?	Include: - MDT referral proforma (attached) - last clinic letter - histology and radiology reports - clinical photographs if available
When?	By the first Friday of the month (for discussion at meeting on the third Friday)

Table 1

Category	For registration only	Full SSMDT discussion
MF Stage I-IIA*	x	
MF Stage I-IIA* refractory to skin-directed therapy		x
MF/SS Stage IIB-IV*		x
Lymphomatoid papulosis	x	
Other pcCD30+ LPD		x
Other CTCL variants		x
CBCL		x
Cutaneous lymphoma suspected but diagnosis unclear		x

*For help with staging, download "Skin Lymphoma" app from Integrated Cancer Research